

# TT 18 Superconductivity - Properties, Electronic Structure, Order Parameter II

Zeit: Montag 10:00–13:00

Raum: TU H2053

## Hauptvortrag

TT 18.1 Mo 10:00 TU H2053

**Two-Gap Superconductivity in MgB<sub>2</sub> — •THOMAS DAHM** — Universität Tübingen, Institut für Theoretische Physik

It is by now generally accepted that the recently discovered superconductor MgB<sub>2</sub> is a superconductor with two energy gaps. In this talk we discuss how this happens and address some of its unusual implications. Measurements of the microwave conductivity on MgB<sub>2</sub> thin films have shown an anomalous coherence peak appearing at significantly lower temperatures than in conventional superconductors [1] and the anisotropy ratio of the upper critical field in MgB<sub>2</sub> single crystals shows an unusual strong temperature dependence. We argue that both of these observations are natural consequences of the existence of the two gaps, if the special Fermi surface structure of MgB<sub>2</sub> is taken into account [1,2]. We also examine consequences for the expected intrinsic nonlinear microwave response in MgB<sub>2</sub> [3].

[1] B.B.Jin, T.Dahm et al. Phys. Rev. Lett. **91**, 127006 (2003).[2] T.Dahm and N.Schopohl, Phys. Rev. Lett. **91**, 017001 (2003).[3] T.Dahm and D.J.Scalapino, Appl. Phys. Lett. **85**, 4436 (2004).

TT 18.2 Mo 10:30 TU H2053

**Interband Pairing Interaction in the Two-Band Superconductor MgB<sub>2</sub> Observed by Tunneling — •J. GEERK<sup>1</sup>, R. SCHNEIDER<sup>1</sup>, G. LINKER<sup>1</sup>, A. ZAITSEV<sup>1</sup>, R. HEID<sup>1</sup>, K.-P. BOHNEN<sup>1</sup>, and H. V. LÖHNESEN<sup>1,2</sup>** — <sup>1</sup>Forschungszentrum Karlsruhe, Institut für Festkörperphysik, P.O.B. 3640, D-76021 Karlsruhe — <sup>2</sup>Physikalisches Institut, Universität Karlsruhe

The anisotropic superconductor MgB<sub>2</sub> was studied by tunneling spectroscopy of tunnel junctions of the sandwich type prepared on as deposited thin films with T<sub>c</sub> values near 32 K. The tunnel junctions revealed an energy gap between 2.5 and 3 meV and phonon induced structures in the tunnelling density of states. The inversion of the tunnel data using the standard single-band Eliashberg equations yielded a so-called effective Eliashberg function with three distinct peaks at 38,58 and 85 meV which is compared to a theoretical calculated counterpart obtained by inversion of superconducting density of states data from two-band Eliashberg equations where electron-phonon spectral functions extracted from ab-initio LDA calculations were inserted. Convincing agreement is found between the experimental and calculated effective Eliashberg functions. Further evaluation reveals that the central peak a 58 meV mainly reflects the shape of the  $\pi$ - $\sigma$  interband pairing interaction which appears in the gap function of the  $\pi$ -sheet in an amplified way due to the large gap of the  $\sigma$ -sheet. It is concluded that the superconductivity on the  $\pi$ -sheet is essentially generated by interband electron-phonon coupling.

TT 18.3 Mo 10:45 TU H2053

**Mixing between ballistic and diffusive motion in superconducting MgB<sub>2</sub> — •MATTHIAS ESCHRIG<sup>1</sup>, KAORI TANAKA<sup>2,3</sup>, DANIEL AGTERBERG<sup>4</sup>, and JUHA KOPU<sup>5</sup>** — <sup>1</sup>Institut für Theoretische Festkörperphysik, Universität Karlsruhe, 76128 Karlsruhe, Germany — <sup>2</sup>Department of Physics and Engineering Physics, University of Saskatchewan, Saskatoon, SK, Canada S7N 5E2 — <sup>3</sup>Argonne National Laboratory, Argonne, IL 60439, U.S.A. — <sup>4</sup>Department of Physics, University of Wisconsin-Milwaukee, P.O. Box 413, Milwaukee, WI 53211, U.S.A — <sup>5</sup>Low Temperature Laboratory, Helsinki University of Technology, PO Box 2200, FIN-02015 HUT

We introduce a model for the two-band superconductor MgB<sub>2</sub> in which the motion of quasiparticles is diffuse in one band (the  $\pi$ -band) and ballistic in the other band (the  $\sigma$ -band). Diffusive and ballistic quasiparticles are coupled by the pairing interaction. We apply this model to examine the electronic structure of vortex cores in MgB<sub>2</sub>. In particular, we study the effects of impurities on the ballistic motion of quasiparticles in the  $\sigma$ -band in the vortex core region under the presence of the hybridization with the diffusive  $\pi$ -band. We find that the induced superconductivity in the  $\pi$  band results in a core size larger than estimated by H<sub>C2</sub> and in weakly bound states, and that those bound states are removed easily by impurities.

TT 18.4 Mo 11:00 TU H2053

**Band filling and interband scattering effects in MgB<sub>2</sub>: C vs Al doping — •JENS KORTUS<sup>1</sup>, O.V. DOLGOV<sup>2</sup>, R.K. KREMER<sup>2</sup>, and A.A. GOLUBOV<sup>3</sup>** — <sup>1</sup>Institut de Physique et Chimie des Matériaux de Strasbourg, 23 rue du Loess, F-67034 Strasbourg Cedex 2, France — <sup>2</sup>MPI-FKF, Heisenbergstr. 1, 70569 Stuttgart — <sup>3</sup>MESA+ Research Institute and Faculty of Science and Technology, University of Twente, 7500 AE Enschede, The Netherlands

We argue, based on band structure calculations and Eliashberg theory, that the observed decrease of T<sub>c</sub> of Al and C doped MgB<sub>2</sub> samples can be understood mainly in terms of a band filling effect due to the electron doping by Al and C. A simple scaling of the electron-phonon coupling constant  $\lambda$  by the variation of the density of states as function of electron doping is sufficient to capture the experimentally observed behavior. Further, we also explain the long standing open question of the experimental observation of a nearly constant  $\pi$  gap as function of doping by a compensation of the effect of band filling and interband scattering. Both effects together generate a nearly constant  $\pi$  gap and shift the merging point of both gaps to higher doping concentrations, resolving the discrepancy between experiment and theoretical predictions based on interband scattering only.

TT 18.5 Mo 11:15 TU H2053

**Density functional theory for superconductors: Applications to MgB<sub>2</sub> and solids under pressure — •ANDREA FLORIS<sup>1</sup>, CESARE FRANCHINI<sup>2</sup>, NEKTARIOS LATHIOTAKIS<sup>1</sup>, GIANNI PROFETA<sup>3</sup>, SANDRO MASSIDDA<sup>2</sup>, and E. K. U. GROSS<sup>1</sup>** — <sup>1</sup>Institut für Theoretische Physik, Freie Universität Berlin, Germany — <sup>2</sup>INFN SLACS, Sardinian Laboratory for Computational Materials Science and Dipartimento di Scienze Fisiche, Università degli Studi di Cagliari, Italy — <sup>3</sup>C. A. S. T. I. - Istituto Nazionale Fisica della Materia (INFN) and Dipartimento di Fisica, Università degli studi dell' Aquila, Italy

Understanding and predicting the properties of superconductors is of both fundamental and technological importance. The discovery of superconductivity in MgB<sub>2</sub>, of its rather high critical temperature (T<sub>c</sub> = 39.5K), and the appearance of multiple gaps, has renewed the interest in conventional superconductivity. Here we present several applications of a novel approach to superconductivity that allows one to calculate material-specific properties, such as the gap and the T<sub>c</sub>, in a truly ab-initio fashion without using any adjustable parameters. Within this approach, we obtained the T<sub>c</sub> and the two gaps of MgB<sub>2</sub> in good agreement with experiment, taking into account the strong anisotropy of both the electron-phonon and the Coulomb interactions. As a further application, we studied the behaviour of T<sub>c</sub> in Li and Al as a function of pressure. Despite their common simple metal structure, these materials show different behaviour upon pressure. While Li undergoes several transitions favouring superconductivity, in Al the electron-phonon coupling decreases with pressure leading to a complete suppression of T<sub>c</sub> around 8GPa.

TT 18.6 Mo 11:30 TU H2053

**Multi-band influence on superconductivity in HoNi<sub>2</sub>B<sub>2</sub>C — •A. WÄLTE<sup>1</sup>, G. FUCHS<sup>1</sup>, YU.G. NAIDYUK<sup>2</sup>, K. NENKOV<sup>1</sup>, S.-L. DRECHSLER<sup>1</sup>, D. SOUPEL<sup>1</sup>, H. ROSNER<sup>3</sup>, J. FREUDENBERGER<sup>1</sup>, K.-H. MÜLLER<sup>1</sup>, G. BEHR<sup>1</sup>, and L. SCHULTZ<sup>1</sup>** — <sup>1</sup>Institut für Festkörper- und Werkstoffforschung Dresden, Helmholtzstr. 20, D-01171 Dresden — <sup>2</sup>B. Verkin Institute for Low Temperature Physics and Engineering, National Academy of Sciences of Ukraine, 47 Lenin Ave., 61103 Kharkiv, Ukraine — <sup>3</sup>MPI CPFS Dresden, Nöthnitzer Str. 40, D-01187 Dresden

Rare-earth nickel borocarbides RNi<sub>2</sub>B<sub>2</sub>C reveal, depending on the rare-earth R, superconductivity, magnetism or even a mixture of these competing phenomena [1]. The reduction of the superconducting ordering temperature T<sub>c</sub> by magnetic pair-breaking fairly follows the Abrikosov-Gorkov model. However, details of the influence of magnetism on the electron system are not yet well understood. From specific heat measurements and point-contact spectroscopy on HoNi<sub>2</sub>B<sub>2</sub>C the phonon density of states F( $\omega$ ) and the spectral function  $\alpha^2F(\omega)$  have been extracted. The characteristic phonon frequency  $\omega_{\text{ph}} \approx 180$  K is similar to  $\omega_{\text{ph}}$  of non-magnetic LuNi<sub>2</sub>B<sub>2</sub>C. Comparing the specific heat jumps of both compounds, an unexpectedly strong deviation from the Abrikosov-Gorkov expectation for HoNi<sub>2</sub>B<sub>2</sub>C is found. A possible explanation is the different influence of pair-breaking on different electron bands.

[1] R.J. Cava *et al.*, Nature 367, January 1994.

TT 18.7 Mo 11:45 TU H2053

**Are intercalated metallochloronitrides electron-phonon mediated superconductors?** — •ROLF HEID and KLAUS-PETER BOHNEN — Institut für Festkörperphysik, Forschungszentrum Karlsruhe

The layered metallochloronitrides XNCl, X=Zr,Hf, have surprised with rather high superconducting transition temperatures of up to 25.5 K after intercalation with Li or Na [1], raising the question to what extent the conventional electron-phonon coupling mechanism is at work. Here we present results of an *ab initio* investigation of the lattice dynamics and electron-phonon coupling of undoped and Li-intercalated ZrNCl applying a density-functional perturbation approach, which allows a calculation of the full momentum dependency of these quantities. The theoretical phonon spectra are found to be in very good agreement with those obtained by neutron scattering experiments [2]. On intercalation, small Fermi surface pockets develop around the K points. This topology gives rise to strongly momentum dependent electron-phonon coupling, which is carried predominantly by two in-plane vibrations of Zr and N. We find that the integrated coupling constant is  $\approx 0.5$ , a value significantly larger than estimated before [3], but still rather small in view of the high  $T_c$  values observed. Implications for the superconducting mechanism will be discussed.

[1] S. Yamanaka *et al.*, Nature **392**, 580 (1998)

[2] P. Adelman *et al.*, J. Low Temp. Phys. **117**, 449 (1999)

[3] R. Weht *et al.*, Europhys. Lett. **48**, 320 (1999)

**Hauptvortrag**

TT 18.8 Mo 12:00 TU H2053

**Point-Contact Spectroscopy on Conventional and Unconventional Superconductors** — •GERNOT GOLL — Physikalisches Institut, Universität Karlsruhe, D-76128 Karlsruhe, Germany

A growing number of metals is found to exhibit exotic types of superconductivity as manifested by their thermodynamic and transport properties. Unconventional superconductivity is observed in many U- and Ce-based heavy-fermion superconductors and oxide superconductors. Point-contact spectroscopy is one of the tools to study the symmetry and nodal structure of the energy gap  $\Delta$ . Andreev reflection of charge carriers at the normal metal/superconductor interface leads to minima at  $V \approx \pm\Delta/e$  in the differential resistance  $dV/dI$  as a function of applied bias  $V$ . In addition, Andreev reflection causes an excess current through the metallic point contact which can be analyzed with respect to the order-parameter symmetry. The talk will review recent developments in this field and focuses on investigations of  $\text{Sr}_2\text{RuO}_4$  as a candidate for *p*-wave superconductivity, and of the heavy-fermion superconductor  $\text{CeCoIn}_5$  as a possible *d*-wave superconductor.

TT 18.9 Mo 12:30 TU H2053

**In-plane angular dependence of the upper critical field of the unconventional superconductor  $\text{CeCoIn}_5$  - Implications for the order parameter symmetry** — •F. WEICKERT<sup>1</sup>, P. GEGENWART<sup>1</sup>, S. HAAS<sup>2</sup>, H. WON<sup>3</sup>, and K. MAKI<sup>2</sup> — <sup>1</sup>Max Planck Institute for Chemical Physics of Solids, Noethnitzer Str. 40, 01187 Dresden, Germany — <sup>2</sup>Department of Physics and Astronomy, University of Southern California, Los Angeles, California 90089-0484, USA — <sup>3</sup>Department of Physics, Hallym University, Chunchon 200-702, South Korea

$\text{CeCoIn}_5$  is a tetragonal heavy fermion system that shows unconventional superconductivity below  $T_c = 2.3\text{K}$  with upper critical fields of about 5T ( $B\parallel c$ ) and 11.5T ( $B\perp c$ ). Below a temperature of 0.7K, the superconducting to normal transition is of first order and at lower temperatures indications for an inhomogeneous Fulde-Ferrell-Larkin-Ovchinnikov state in close vicinity to  $B_{c2}(T)$  have been discovered.

Here we use resistivity measurements to study the in-plane angular dependence of the upper critical field  $B_{c2}(\theta)$  at 0.1K. A fourfold oscillation with 2% amplitude is observed. Since  $B_{c2}$  for fields parallel to the [110] direction is larger than along the [100] direction, our results would be compatible with a  $d_{xy}$  superconducting order parameter.

TT 18.10 Mo 12:45 TU H2053

**Thermoelectric power of  $\text{V}_3\text{Si}$  single crystal near the structural transition** — •JUN SUNG KIM and REINHARD K KREMER — MPI-FKF, 70569 Stuttgart

$\text{V}_3\text{Si}$ , one of the A-15 superconductors, has attracted a lot of interest in experimental and theoretical studies because of the relatively high superconducting transition temperature and the unclear role of the lattice

instability. The structural phase transition from cubic to tetragonal occurs at  $T_M \sim 23\text{K}$ , just above the superconducting transition  $T_C = 16.7\text{K}$ . To study the implication of the structural phase transition on the electronic properties, we have carried out highly temperature resolved thermoelectric power measurements in the regime of the structural transition on a single crystal which according to heat capacity measurements shows a clear transition at  $T_M$ . Above  $T_M$ , the thermoelectric power ( $S$ ) shows a nonlinear temperature dependence with a knee structure near  $T \sim 100\text{K}$ , which is consistent with previous measurements on polycrystalline samples.[1] However, it is found that there is a clear slope change in  $S(T)$  near  $T_M$  on the single crystal, and  $S(T)$  rapidly decreases below  $T_M$ . These results indicate that the scattering asymmetry between hole and electron is reduced due to the structural phase transition. The change of the density of states near the Fermi level and electron-phonon coupling will be discussed as an origin of temperature dependence of  $S(T)$ .

[1] M. P. Sarachik, *et al.* Can. J. Phys. 41, 1542 (1963).