

Symposium Soft X-ray induced ultrafast processes on the atomic scale (SYSX)

gemeinsam veranstaltet
vom Fachverband Atomphysik und
vom Fachverband Molekülphysik

<p>Thomas Möller Technische Universität Berlin Institut für Optik und Atomare Physik Hardenbergstrasse 36 10623 Berlin thomas.moeller@physik.tu-berlin.de</p>	<p>Christian Spielmann Physikalisches Institut EP1 Universität Würzburg Am Hubland D-97074 Würzburg spielmann@physik.uni-wuerzburg.de</p>
---	---

Übersicht der Hauptvorträge und Fachsitzungen

(Hörsaal 6G)

Hauptvorträge

SYSX 1.1	Do	14:00–14:30	6G	Unraveling ultra fast dynamic processes at selected atomic sites — •ALEXANDER FOEHLISCH
SYSX 1.2	Do	14:30–15:00	6G	Ultrafast processes on the atomic scale studied with hard (and soft) x-radiation — •CHRISTIAN BRESSLER
SYSX 1.3	Do	15:00–15:30	6G	Interaction of clusters with intense soft x-ray radiation from the FLASH VUV-FEL — •CHRISTOPH BOSTEDT, HEIKO THOMAS, EKATERINA EREMINA, MATTHIAS HOENER, HUBERTUS WABNITZ, TIM LAARMANN, RUBENS DE CASTRO, THOMAS MÖLLER
SYSX 1.4	Do	15:30–16:00	6G	Elektronendynamik bei der Erzeugung von XUV-Attosekundenpulsen — •MILUTIN KOVACEV
SYSX 2.1	Do	16:30–17:00	6G	Atomic photoionization by femtosecond soft X-ray pulses — •MATHIAS RICHTER, SERGEY V. BOBASHEV, ANDREI A. SOROKIN, KAI TIEDTKE, HUBERTUS WABNITZ, MICHAEL WELHÖFER

Fachsitzungen

SYSX 1.1–1.4	Do	14:00–16:00	6G	Soft X-ray induced ultrafast processes I
SYSX 2.1–2.3	Do	16:30–17:30	6G	Soft X-ray induced ultrafast processes II

SYSX 1: Soft X-ray induced ultrafast processes I

Zeit: Donnerstag 14:00–16:00

Raum: 6G

Hauptvortrag SYSX 1.1 Do 14:00 6G
Unraveling ultra fast dynamic processes at selected atomic sites — ●ALEXANDER FOEHLISCH — Department Physik, Universität Hamburg, Luruper Chaussee 149, 22761 Hamburg

Ultra fast electron and nuclear dynamics play a decisive role in processes like photo- and electrochemistry as well as in molecular electronics and single electron devices. If we are interested in a dynamic process within a complex systems it is of equal importance to address the temporal evolution of the wave packet and the question at which atomic center the dynamic has started. These requirements can be matched with the element specific information from soft x-ray spectroscopy [1,2]. The novel X-ray source of the Free-Electron-Laser provides us with an unprecedented peak brilliance of femtosecond X-ray pulses. With these femtosecond radiation pulses at the Free Electron Laser in Hamburg (FLASH) we have investigated ultra fast dynamic processes in a pump-probe set-up in combination with an optical short-pulse laser.

[1] A. Föhlisch, P. Feulner, F. Hennies, A. Fink, D. Menzel, D. Sanchez-Portal, P. M. Echenique, W. Wurth, Direct observation of electron dynamics in the attosecond domain. *Nature* 436, 373 (2005).

[2] S. Vijayalakshmi, A. Föhlisch, F. Hennies, A. Pietzsch, M. Nagasono, W. Wurth. Surface projected electronic band structure and Adsorbate charge transfer dynamics: Ar adsorbed on Cu(111) and Cu(100). *Chem. Phys. Lett.* 427, 91-95 (2006).

Hauptvortrag SYSX 1.2 Do 14:30 6G
Ultrafast processes on the atomic scale studied with hard (and soft) x-radiation — ●CHRISTIAN BRESSLER — LSU-ISIC-BSP. EPFL. CH-1015 Lausanne, Switzerland

Electronic structure changes are at the origin of chemical reactivity, which drive the forming and breaking of bonds. Ultrafast x-ray absorption fine structure spectroscopy (XAFS) delivers both the electronic and geometric transient structure changes, when interfaced with a femtosecond laser in a pump-probe scheme. In addition, XAFS methods are element-selective and can be applied to disordered bulk systems, in particular liquids. On the femtosecond time scale one would obtain a rather complete picture via XAFS, since the propagation from a reaction center is limited to a few Angstroms on these time scales. We have recently recorded high-quality transient XAFS spectra at 3rd generation synchrotrons with currently < 100 ps temporal resolution, corresponding to the electron bunch width at the synchrotron. Examples of excited state structures of solvated coordination chemistry compounds and of short-lived atomic radicals in aqueous solution will be presented. Extending these studies into the femtosecond time domain is straightforward in concept, while pico- to nanosecond resolved studies still offer a high potential for research on short-lived reaction intermediates, whose structures can not be determined by other means. The continued efforts using pulsed x-rays on the femto- to nanosecond time scales will be closely linked to the developments of novel photon sources, including, e.g., femtosecond x-rays at SLS and BESSY, next to soft x-rays < 1 keV to become available at FLASH.

Hauptvortrag SYSX 1.3 Do 15:00 6G
Interaction of clusters with intense soft x-ray radiation from the FLASH VUV-FEL — ●CHRISTOPH BOSTEDT¹, HEIKO THOMAS¹, EKATERINA EREMINA¹, MATTHIAS HOENER¹, HUBERTUS WABNITZ², TIM LAARMANN³, RUBENS DE CASTRO⁴, and THOMAS MÖLLER¹ — ¹TU Berlin, Germany — ²DESY, Hamburg, Germany — ³MBI, Berlin, Germany — ⁴LNLS, Campinas, Brazil

The interaction of rare gas clusters with intense vacuum ultraviolet radiation from the DESY free electron lasers has yielded many surprising results. In first experiments with 100 nm radiation unexpected high energy absorption involving plasma heating processes was measured and new theoretical explanations were suggested. The new DESY FLASH FEL delivers radiation down to first 32 and later also 13 nm wavelengths with intensities up to 10^{14} W/cm². We have performed first experiments with a combined time-of-flight spectroscopy and imaging approach. The new data give evidence that the dominant physical drivers in the light - matter interaction at reduced wavelength change and plasma heating is only of minor importance. At 32 nm the ionic fragments from the exploding clusters are ejected with only 25 eV and the electrons are emitted from the clusters serially. The atoms within the cluster move by only 3 Angstrom during the pulse duration making imaging experiments possible. At 13 nm wavelength first experiments about the ionization dynamics including core levels could be performed with Xe clusters. These measurements give some hints that the energy absorption involving core levels in the cluster may be reduced compared to atoms.

Hauptvortrag SYSX 1.4 Do 15:30 6G
Elektronendynamik bei der Erzeugung von XUV-Attosekundenpulsen — ●MILUTIN KOVACEV — Institut für Quantenoptik, Leibniz Universität Hannover

Die Erforschung ultrakurzer, intensiver Laserpulse und deren Wechselwirkung mit Materie ist ein hochaktuelles Forschungsgebiet. Eine Vielzahl faszinierender Effekte und Anwendungen konnten in den letzten Jahren verwirklicht werden.

Insbesondere die Erzeugung hoher optischer Harmonischer im weichen Röntgenbereich ist durch eine hohe Brillanz (10^{27} Phot/Puls/mm²/mrad²/0,1%BW), exzellente Kohärenzeigenschaften und eine ultrakurze Pulsdauer gekennzeichnet. Darüber hinaus sind Laser-Erzeugte XUV Strahlungsquellen kompakt und kostengünstig, welches deren vielfältigen Einsatz in Laboratorien begünstigt.

Das Harmonischen-Spektrum ermöglicht durch seine große Bandbreite die Erzeugung ultrakurzer Pulse im Attosekundenbereich ($1\text{as} = 10^{-18}\text{s}$). Der Erzeugungsprozess dieser Pulse unterliegt einer anspruchsvollen Elektronendynamik, auf die in diesem Beitrag eingegangen wird. Die Kontrolle der Elektronendynamik und insbesondere auch die Synchronisation des Phasenganges der Harmonischen sind für Anwendungen in der Attosekundenphysik entscheidend und sollen hier diskutiert werden.

SYSX 2: Soft X-ray induced ultrafast processes II

Zeit: Donnerstag 16:30–17:30

Raum: 6G

Hauptvortrag SYSX 2.1 Do 16:30 6G
Atomic photoionization by femtosecond soft X-ray pulses — ●MATHIAS RICHTER¹, SERGEY V. BOBASHEV², ANDREI A. SOROKIN^{1,2}, KAI TIEDTKE³, HUBERTUS WABNITZ³, and MICHAEL WELHÖFER⁴ — ¹Physikalisch-Technische Bundesanstalt, PTB, Abbestraße 2-12, D-10587 Berlin, Germany — ²Ioffe Physico-Technical Institute, Polytekhnicheskaya 26, 194021 St. Petersburg, Russia — ³Deutsches Elektronen-Synchrotron, DESY, Notkestraße 85, D-22603 Hamburg, Germany — ⁴Universität Hamburg, Luruper Chaussee 149, 22761 Hamburg, Germany

The contribution refers to recent results on atomic photoionization obtained in focused photon beams at the new free-electron laser (FEL) in Hamburg (FLASH) by ion time-of-flight mass/charge spectroscopy. The experiments were performed with emphasis on a quantitative de-

termination of pulse energy in the order of 10 microjoule and focus diameter down to 3 micrometer which yields, together with the estimated pulse duration of 10 to 30 fs, irradiance levels of up to $1\text{e}16$ W/cm² achieved. Strong non-linear behavior due to multi-photon processes was observed on helium, neon, and xenon atoms studied as a function of photon intensity. Results are discussed with regard to the role of sequential and direct multi-photon ionization and a theoretical description by perturbative and non-perturbative methods.

Hauptvortrag SYSX 2.2 Do 17:00 6G
Photoelektronenspektroskopie von Atomen und Molekülen nach Multiphoton-Ionisation durch FEL-Strahlung — ●M. BRAUNE¹, A. REINKÖSTER¹, J. VIEFHAUS², B. LOHMANN³ und U. BECKER¹ — ¹Fritz-Haber-Institut der MPG, 14195 Berlin — ²DESY,

22607 Hamburg — ³Universität Münster, 48149 Münster

Photoelektronenspektroskopie von Atomen und Molekülen in der Gasphase ist eine weit verbreitete Methode für die Erforschung der elektronischen Struktur der Materie. Mit der Inbetriebnahme des VUV Freielektronen-Lasers FLASH in Hamburg wurden erstmals Multiphoton-Ionisationsexperimente möglich, wobei Ionisation durch mehrere Photonen sowohl simultan als auch sequentiell auftritt. In Experimenten an Edelgasen war die Intensität der simultanen Mehrfach-Ionisation um zwei Größenordnungen kleiner die der Einfach-Ionisation der neutralen Atome, die Signale der sequentiellen Zweifach-Ionisation erreichten jedoch die gleiche Größenordnung wie die der Einfach-Ionisation. Innerhalb der Dauer eines einzelnen FEL-Lichtpulses können also ionische Targets beträchtlicher Dichte erzeugt und anschließend weiter ionisiert werden. Das eröffnet die Möglichkeit, mittels winkelauflösender und spinsensitiver Photoelektronenspektroskopie die verschiedenen Ionisationsstufen gasförmiger Materie zu studieren und dabei insbesondere das Zusammenspiel von sequentieller und simultaner Ionisation in Multiphotonen-Prozessen zu untersuchen. Erste Ergebnisse von winkelauflösenden Flugzeitmessungen von Photoelektronen verschiedener Gas-Targets werden vorgestellt.

SYSX 2.3 Do 17:15 6G

VUV FEL photodissociation imaging of HeH⁺ at 32nm
— •SIMON ALTEVOGT¹, HENRIK PEDERSEN¹, BRANDON JORDON-THADEN¹, ODED HEBER², MICHAEL RAPPAPORT², DIRK SCHWALM¹, DANIEL ZAJFMAN², JOACHIM ULLRICH¹, ANDREAS WOLF¹, ROLF TREUSCH³, NATALIA GUERASSIMOVA³, MICHAEL MARTINS⁴, JON HOEFT⁴, and MICHAEL WELHÖFER⁴ — ¹Max-Planck-Institut für Kernphysik, 69117 Heidelberg, Germany — ²Department of Particle Physics, Weizman Institute of Science, Rehovot, 76100, Israel — ³HASYLAB, DESY, 22607 Hamburg, Germany — ⁴Institut für Experimentalphysik, Universität Hamburg, 22671 Hamburg, Germany

Many molecules are most efficiently photofragmented by VUV photons (100 – 30nm). With fast ion beams crossed with an intense photon beam the momentum of neutral fragments after photodissociation reactions can be measured accurately event by event. Such experiments have become possible with the VUV free-electron laser FLASH at DESY. Using the new ion infrastructure TIFF the kinetic energy and angular distributions of neutral He products was measured for HeH⁺ photofragmentation at 32nm together with the absolute cross section for this channel. Excited levels of ¹Π symmetry, not so far considered in photodissociation calculations, are found to dominate the process.