

Intersectional Symposium Attosecond Physics: Atoms, Molecules and Condensed Matter (SYAP)

lead by the Surface Science Division (O)

Martin Wolf
Fritz-Haber-Institut der MPG
Berlin

Ferenc Krausz
MPI für Quantenoptik
LMU München

Ralph Ernstorfer
Fritz-Haber-Institut der MPG
Berlin

Within the last decade, attosecond physics has established itself as a new field of research. Attoseconds are the time scale of fundamental electronic effects in atoms, molecules as well as in solids. In recent years, new spectroscopic techniques were evolving in a series of proof-of-principle experiments probing attosecond electron dynamics in the time domain. These new experimental findings challenge theoretical approaches by invalidating common approximations. For instance, it is an open question to which extent the concept of quasi-particles or a band structure can be applied on the intrinsic time scales of electron-electron correlations. This symposium aims to provide interdisciplinary perspectives on attosecond phenomena and to stimulate discussions on relevant theoretical concepts in atomic, molecular and condensed matter physics.

Overview of Invited Talks and Sessions

(lecture room HSZ 01)

Invited Talks

SYAP 1.1	Mon	10:30–11:00	HSZ 01	Observing Intra-atomic Electron Correlation by Tunnelling and Recollision — ●PAUL CORKUM
SYAP 1.2	Mon	11:00–11:30	HSZ 01	Attosecond time-resolved molecular electron dynamics — ●MARC VRACKING
SYAP 1.3	Mon	11:30–12:00	HSZ 01	Opportunities in Attosecond Science using Core Level Spectroscopy — ●ANDERS NILSSON
SYAP 1.4	Mon	12:00–12:30	HSZ 01	Attosecond spectroscopy on solid surfaces — ●REINHARD KIENBERGER
SYAP 1.5	Mon	12:30–13:00	HSZ 01	Condensed matter effects in attosecond physics — ●PEDRO M. ECHENIQUE

Sessions

SYAP 1.1–1.5	Mon	10:30–13:00	HSZ 01	Attosecond Physics: Atoms, Molecules, Condensed Matter
--------------	-----	-------------	--------	---

SYAP 1: Attosecond Physics: Atoms, Molecules, Condensed Matter

Time: Monday 10:30–13:00

Location: HSZ 01

Invited Talk SYAP 1.1 Mon 10:30 HSZ 01
Observing Intra-atomic Electron Correlation by Tunnelling and Re-collision — ●PAUL CORKUM — Joint Attosecond Science Lab, 100 Sussex Drive, Ottawa, ON, Canada

Laser induced tunnelling imposes two filters on a quantum system * a directional filter in momentum space and an ionization potential filter. Together they allow us to preferentially address specific orbitals or to launch valence electron wave packets in the ion. Using sequential ionization, we show how circular polarized light can image electron correlations in rare gas atoms and HCl [1]

In linear polarized light the tunnelled electron re-collides. During the re-collision it can probe its correlated hole through high harmonic generation [2]. If the hole moves the dynamics shows up in the structure of the high harmonic spectrum. We show how attosecond dynamics can be observed experimentally.

[1] A. Fleischer, H. Wörner, D.M. Villeneuve, A. Staudte and P. B. Corkum, *Probing Electron Correlations by Laser-Induced Tunnel Ionization* submitted for publication.

[2] O. Smirnova, Y. Mairesse, S. Patchkovskii, N. Dudovich, D. M. Villeneuve, P. B. Corkum, and M. Y. Ivanov, *Probing Inner Orbitals and Electron Tunnelling in Molecules with High Harmonic Interferometer*, Nature, 460, 972 (2009).

Invited Talk SYAP 1.2 Mon 11:00 HSZ 01
Attosecond time-resolved molecular electron dynamics — ●MARC VRAKKING — Max-Born-Institut, Berlin

Using attosecond light sources based on high-harmonic generation (HHG), pump-probe experiments can be performed where electron dynamics is studied on its natural timescale, providing insight into the fundamental role that electrons play in photo-induced processes.

In my talk I will present some of the first applications of these techniques in molecular science, focusing on two-color experiments where several small molecules were exposed to a sequence of one or more attosecond pulses and an infrared field. In addition to the attosecond time-structure accompanying HHG, the short wavelengths in HHG allow to use ejected photoelectrons as a probe of the (time-evolving) molecular structure, which amounts to *illuminating the molecule from within*. I will present the present status of our experiments on this topic.

Invited Talk SYAP 1.3 Mon 11:30 HSZ 01
Opportunities in Attosecond Science using Core Level Spectroscopy — ●ANDERS NILSSON — SLAC National Accelerator Laboratory, Stanford University, USA

I will give an overview how core level spectroscopy can be used to probe ultrafast processes on surfaces and in the liquid phase. In the frequency domain, we observe electronic transfer and nuclear motions of hydrogen atoms in the decay spectra based on the concept of using the life time of the inner shell hole as an internal clock. This has been used to follow dissociation events in gas phase molecules, electron transfer processes of adsorbates on surfaces, in liquid water and ions in aqueous systems. There are new opportunities with x-ray free electron lasers, such as LCLS, where x-ray pulses enter into attosecond regime opening up for new class of studies that will be discussed.

Invited Talk SYAP 1.4 Mon 12:00 HSZ 01
Attosecond spectroscopy on solid surfaces — ●REINHARD KIENBERGER — Fakultät für Physik, TUM, James Franck Str., 85748 Garching

The generation of ever shorter pulses is a key to exploring the dynamic behavior of matter on ever shorter time scales. Attosecond XUV pulses together with the few-cycle (few-femtosecond) IR laser pulses used for their generation have opened the way to the development of a technique for attosecond sampling of electrons ejected from atoms or molecules or solids. Atoms exposed to a few oscillation cycles of intense visible or near-infrared light are able to emit a single electron and XUV photon wavepacket of sub-femtosecond duration [1]. After the full characterization of these tools, first experiments have been carried out to measure sub-femtosecond behavior of matter. Recently, the dynamics of the photoionization process on solids has been studied [2]. Not only that attosecond metrology now enables clocking on surface dynamics, but also the individual behaviour of electrons of different type (core electrons vs. conduction band electrons) can be resolved. Here, we measured a time delay of about 100 as on the emission of the aforementioned two types of electrons. For absolute determination of the emission process, the *travel* time of electrons through defined numbers of adlayers is investigated.

[1]*R. Kienberger et al., Nature 2004, 427, 817. [2]*A. Cavalieri et al., Nature 2007, 449, 1029.

Invited Talk SYAP 1.5 Mon 12:30 HSZ 01
Condensed matter effects in attosecond physics — ●PEDRO M. ECHENIQUE — Donostia International Physics Center (DIPC) and Unidad de Física de Materiales CSIC-UPV/EHU, Manuel de Lardizabal Pasealekua, E-20018 Donostia, Basque Country, Spain

Recent theoretical and experimental spectroscopic experimental results will be discussed with an emphasis on the relative importance of atomic versus condensed matter physics.