Symposium Frontiers of Nanomechanics (SYNM)

jointly organized by the Low Temperature Physics Division (TT), the Dynamics and Statical Physics Division (DY), and the Semiconductor Physics Division (HL)

Florian Marquardt Universität Erlangen-Nürnberg Institut für Theoretische Physik Staudtstr. 7 D-91058 Erlangen Florian.Marquardt@physik.unierlangen.de Milena Grifoni Institut I - Theoretische Physik Universität Regensburg D-93040 Regensburg milena.grifoni@physik.uniregensburg.de Felix von Oppen Fachbereich Physik Freie Universität Berlin Arnimallee 14 D-14195 Berlin vonoppen@physik.fu-berlin.de

Nanomechanical systems are receiving strong attention because of their potential both in terms of fundamental quantum physics and for applications such as ultrasensitive detection of small masses, forces and displacements, and as a method for transferring and manipulating quantum information. This symposium highlights recent experimental results at the forefront of international research, including coupling to a superconducting qubit, to microwave resonators, and to the optical radiation field.

Overview of Invited Talks and Sessions

(lecture room H 0105)

Invited Talks

SYNM 1.1	Wed	15:00 - 15:30	H 0105	Mechanical resonators in the quantum regime — •ANDREW N. CLE- LAND
SYNM 1.2	Wed	15:30 - 16:00	H 0105	Quantum optomechanics: exploring the interface between quantum
SYNM 1.3	Wed	16:00-16:30	H 0105	Integrated transduction and coherent control of high Q nanome-
	XX 7 1	16 00 15 00		chanical systems using dielectric gradient forces $-\bullet$ Eva M. Weig
SYNM 1.4 SYNM 1.5	Wed Wed	16:30-17:00 17:00-17:30	H 0105 H 0105	Optomechanical crystals — •Oskar Painter

Sessions

SYNM 1.1–1.5 Wed 15:00–17:30 H 0105 Frontiers of Nanomechanics

SYNM 1: Frontiers of Nanomechanics

Time: Wednesday 15:00-17:30

Invited TalkSYNM 1.1Wed 15:00H 0105Mechanical resonators in the quantum regime- • ANDREW N.CLELAND— Department of Physics, University of California, SantaBarbara CA 93106 USA

I will describe our recent experiments, representing about ten years' development of nanomechanical and quantum circuit technology, which culminated in our formulating and creating a quantum mechanical resonator that could *easily* be prepared in quantum (non-classical) states of mechanical vibration. Key requirements included a mechanical design that supported a microwave-frequency mechanical resonance; using a piezoelectric material in order to achieve very strong electromechanical coupling; and employing a Josephson junction, implemented as a phase quantum bit (qubit), to measure and interact with the mechanical resonator. Operating at 25 mK on the mixing chamber of a dilution refrigerator, this integrated electromechanical system can be cooled to its quantum ground state without additional intervention. Then, employing the extraordinary nonlinearity provided by the Josephson qubit, and the coherent interactions of this qubit and the mechanical resonator, we were able to prepare and measure nonclassical mechanical states of motion in the resonator.

Invited TalkSYNM 1.2Wed 15:30H 0105Quantum optomechanics:exploring the interface betweenquantum physics and gravity- •MARKUS ASPELMEYERVienna Center for Quantum Science and Technology (VCQ), Faculty ofPhysics, University of Vienna, Austria

Quantum optics provides a high-precision toolbox to enter and to control the quantum regime of the motion of massive mechanical objects. This opens the door to a hitherto untested parameter regime of macroscopic quantum physics. Due to the large available mass range – from picograms in nanomechanical waveguides to kilograms in mirrors for gravitational wave detection – it becomes possible to explore the fascinating interface between quantum physics and (quantum) gravity in table-top quantum optics experiments. I will discuss a few examples.

Invited TalkSYNM 1.3Wed 16:00H 0105Integrated transduction and coherent control of high Qnanomechanical systems using dielectric gradient forces•Eva M. Weig — LMU München

Nanomechanical resonators, freely suspended bridges with nanoscale cross-sections, are receiving an increasing amount of attention both in fundamental experiments and sensing applications. In particular, prestressed silicon nitride is explored as a high Q material for nanomechanical systems (NEMS): As a consequence of the intrinsic tensile stress of the nitride film, room temperature quality factors of several 100,000 are observed in the 10 MHz eigenfrequency range. To take advantage of the large quality factor, non-dissipative transduction and manipulation schemes have to be developed which do not introduce extra damping.

We employ optically or electrically induced gradient forces to implement dielectric transduction as an efficient way to actuate and probe high Q nanomechanical resonators and to control the resonator eigenfrequency. The displacement sensitivity can be enhanced considerably by taking advantage of near-field coupling to a cavity, which, at the same time allows to benefit from back-action induced dynamics, enabling to cool the Brownian motion or to initiate cavity-pumped self-oscillation. A realization based on a room temperature microwave cavity offers a plug and play solution for all-integrated transduction, and can be employed to probe the dynamics of two strongly coupled resonator modes.

Invited Talk SYNM 1.4 Wed 16:30 H 0105 Cavity optomechanics with microwave photons — •JOHN TEUFEL — NIST, Boulder, USA

Accessing the full quantum nature of a macroscopic mechanical oscillator first requires elimination of its classical, thermal motion. The flourishing field of cavity optomechanics provides a nearly ideal architecture for both preparation and detection of mechanical motion at the quantum level. We realize a microwave cavity optomechanical system by coupling the motion of an aluminum membrane to the resonance frequency of a superconducting circuit [1]. By exciting the microwave circuit below its resonance frequency, we damp and cool the membrane motion with radiation pressure forces, analogous to laser cooling of the motion of trapped ions. The microwave excitation serves not only to cool, but also to monitor the displacement of the membrane. A nearly shot-noise limited, Josephson parametric amplifier is used to detect the mechanical sidebands of this microwave excitation and quantify the thermal motion as it is cooled with radiation pressure forces to its quantum ground state [2].

[1] Teufel, J. D. et al., Nature 471, 204-208 (2011).

[2] Teufel, J. D. et al., Nature 475, 359-363 (2011).

Invited Talk SYNM 1.5 Wed 17:00 H 0105 Optomechanical crystals — •OSKAR PAINTER — Thomas J. Watson, Sr., Laboratory of Applied Physics, California Institute of Technology, Pasadena CA 91125 USA

In the last several years, rapid advances have been made in the field of cavity optomechanics, in which the usually feeble radiation pressure force of light is used to manipulate (and precisely monitor) mechanical motion. These advances have moved the field from the multi-km interferometer of a gravitational wave observatory, to the optical table top, and now all the way down to a silicon microchip. In this talk I will describe these advances, and discuss our own work to realize radiation pressure within nanoscale structures in the form of coupled photonic and phononic crystals (dubbed optomechanical crystals). Applications of these new nano-opto-mechanical systems include: all-optically tunable photonics, optically powered RF and microwave oscillators, and precision force/acceleration and mass sensing. Additionally there is the potential for these systems to be used in hybrid quantum networks, enabling storage or transfer of quantum information between disparate quantum systems. I will introduce several conceptual ideas regarding phonon-photon translation and slow light effects which may be used in such quantum settings, and discuss recent experiments to realize them in practice.

Location: H 0105