

TT 14: Matter At Low Temperature: Multiferroics (jointly with MA, DF, DS, KR)

Time: Monday 16:45–18:00

Location: H 3005

Invited Talk

TT 14.1 Mon 16:45 H 3005

Multiferroicity in an organic charge-transfer salt: Electric-dipole-driven magnetism — •PETER LUNKENHEIMER¹, JENS MÜLLER², STEPHAN KROHNS¹, FLORIAN SCHRETTLE¹, ALOIS LOIDL¹, and MICHAEL LANG² — ¹Experimental Physics V, Center for Electronic Correlations and Magnetism, University of Augsburg, Germany — ²Institute of Physics, Goethe-University Frankfurt, Germany

Multiferroics, showing simultaneous electrical and magnetic ordering, are remarkable materials from both an academic and technological point of view. A prominent mechanism is the spin-driven ferroelectricity, often found in materials with helical spin order. However, recently a different mechanism, namely purely electronic ferroelectricity, where charge order breaks inversion symmetry, has attracted considerable interest. In the present talk, I will treat examples for both types of multiferroics like perovskite manganites and magnetite, concentrating on their dielectric properties, which often are only poorly characterized. An especially interesting case is a two-dimensional organic charge-transfer salt, which shows ferroelectricity, accompanied by antiferromagnetic spin order and belongs to a new class of multiferroics [1]. In this material, the ferroelectric ordering leads to a breaking of spin frustration, which triggers simultaneous dipolar and spin order. Hence, here the spin order is driven by the ferroelectricity, in marked contrast to the spin-driven ferroelectricity in helical magnets.

[1] P. Lunkenheimer, J. Müller, S. Krohns, F. Schrettle, A. Loidl, B. Hartmann, R. Rommel, M. de Souza, C. Hotta, J.A. Schlueter, M. Lang, preprint (arXiv:1111.2752).

TT 14.2 Mon 17:15 H 3005

Critical dynamics in LiCuVO₄ — •CHRISTOPH GRAMS¹, MAXIMILIAN SCHALENBACH¹, DANIEL NIERMANN¹, SANDRA NIESEN¹, PETRA BECKER², and JOACHIM HEMBERGER¹ — ¹II. Physikalisches Institut, Universität zu Köln, Germany — ²Institut für Kristallographie, Universität zu Köln, Germany

Without an external magnetic field LiCuVO₄ has a phase transition into a cycloidal spin ordered phase below 2.3 K where it simultaneously is antiferromagnetic and ferroelectric. The transition temperature of this phase transition can be lowered with increasing magnetic field.

Ferroelectric phase transitions are of continuous fashion and are accompanied with a symmetry lowering that yields soft modes. Near the critical point the dynamics of the ionic polarization mechanisms are slowed down and therefore denoted as critical (“critical slowing down”).

We observe the critical dynamics of the low temperature multiferroic

phase transition of LiCuVO₄ with broadband dielectric spectroscopy. Therefore the compound’s dielectric response to an external electric AC field of frequencies from 10 mHz to several GHz in the temperature range 20 mK to 300 K was measured in dependence of magnetic fields up to 14 T.

Work supported by the DFG through SFB 608.

TT 14.3 Mon 17:30 H 3005

Vortex domain walls in helical magnets — •THOMAS NATTERMANN¹, FUXIAN LI², and VALERY L. POKROVSKY^{2,3} — ¹Institut für Theoretische Physik, Universität zu Köln, D-50937 Köln, Germany — ²Department of Physics, Texas A&M University, College Station, Texas 77843-4242 — ³Landau Institute for Theoretical Physics, Chernogolovka, Moscow District, 142432, Russia

The theory of domain walls in both centrosymmetric and non-centrosymmetric helical magnets is presented. With the exception of discrete orientations domain walls consist of an array of parallel vortex lines, their width is only weakly depending anisotropy, in contrast to ferromagnets and antiferromagnets. In conical phases vortex walls carry Berry phase flux which gives rise to an anomalous Hall effect. In multi-ferroics vortices are electrically charged.

TT 14.4 Mon 17:45 H 3005

Infrared and THz spectroscopy in multiferroic Eu_{1-x}Ho_xMnO₃ — •ZHENYU CHEN¹, MICHAEL SCHMIDT¹, FRANZ MAYR¹, ZHE WANG¹, A.A. MUKHIN², JOACHIM DEISENHOFER¹, and ALOIS LOIDL¹ — ¹Experimentalphysik V, EKM, University of Augsburg, 86135 Augsburg, Germany — ²General Physics Institute of the Russian Academy of Sciences, 119991 Moscow, Russia

We investigated Eu_{1-x}Ho_xMnO₃ with the concentration varying from 0.1 to 0.5. On cooling, Eu_{0.9}Ho_{0.1}MnO₃ enters an incommensurate antiferromagnetic phase, which turns into a commensurate antiferromagnetic one at lower temperatures. Doping leads to ferroelectricity with polarization parallel to the a-axis, which flips to P//c by further doping. In order to detect the coupling between low energy phonons and electromagnons [1-4], we performed systematic polarization dependent IR and THz studies. The data will be compared to TbMnO₃ and the related system Eu_{1-x}Y_xMnO₃.

[1] A.Pimenov et al., Nature Phys., 2, 97(2006).

[2] A.Pimenov et al., Phys. Rev. B, 77, 014438(2008).

[3] N.Kida et al., Phys. Rev. B, 78, 104414(2008).

[4] R.Valdés Aguilar et al., Phys. Rev. Lett., 102, 047203(2009).