

SYQG 2: Quantum meets gravity and metrology II

Time: Tuesday 14:30–16:00

Location: E 415

SYQG 2.1 Tue 14:30 E 415

Differences between neutron and atom interferometry — ●ENNO GIESE¹, DANIEL M. GREENBERGER², ERNST M. RASEL³, and WOLFGANG P. SCHLEICH¹ — ¹Institut für Quantenphysik and Center for Integrated Quantum Science and Technology (IQST), Universität Ulm, D-89069 Ulm, Germany. — ²The City College of New York, 160 Convent Ave, New York, NY 10031, USA. — ³Institut für Quantenoptik, Leibniz Universität Hannover, Welfengarten 1, D-30167 Hannover, Germany.

Triggered by the controversy whether atom interferometers are sensitive to relativistic effects such as the red shift, the difference between neutron and atom interferometers drew our attention. In contrast to a the first naive guess, subtle distinctions can be found. In general, the difference between these two types of interferometers can be traced back to the different scattering processes. A conventional neutron interferometer uses crystals as beam splitters and mirrors. This scattering mechanism can be understood in terms of conventional Bragg diffraction.

In contrast to that, a variety of scattering mechanisms can be applied to atom interferometers. In this talk, we focus on atomic Bragg scattering, where the atoms interact with a standing light wave and change just their external degree of freedom. We show that by a careful arrangement and tuning of the lasers this diffraction process can be changed so that a neutron interferometer is mimicked. The phases accumulated along both paths are different in comparison to the usual atom interferometer which leads to a measurable phase shift.

Group Report

SYQG 2.2 Tue 14:45 E 415

A miniaturized, high flux BEC source for precision interferometry — ●WALDEMAR HERR¹, ERNST MARIA RASEL¹, and THE QUANTUS TEAM^{1,2,3,4,5,6,7,8,9} — ¹Institut für Quantenoptik, Leibniz Universität Hannover — ²ZARM, Universität Bremen — ³Institut für Physik, HU Berlin — ⁴Institut für Laser-Physik, Universität Hamburg — ⁵Institut für Quantenphysik, Universität Ulm — ⁶Institut für angewandte Physik, TU Darmstadt — ⁷MUARC, University of Birmingham — ⁸FBH, Berlin — ⁹MPQ, Garching

Atom chips have proven to be excellent sources for the fast production of ultra-cold gases due to their outstanding performance in evaporative cooling. However, the total number of atoms has previously been limited by the small volume of their magnetic traps. To overcome this restriction, we have developed a novel loading scheme that allows us to produce Bose-Einstein condensates of a few 10^5 ⁸⁷Rb atoms every two seconds. The apparatus is designed to be operated in microgravity at the drop tower in Bremen, where even higher numbers of atoms can be achieved in the absence of any gravitational sag.

Using the drop tower's catapult mode, our setup will perform atom interferometry during nine seconds in free fall. Thus, the fast loading scheme allows for interferometer sequences of up to seven seconds – interrogation times which are inaccessible for ground based devices.

The QUANTUS project is supported by the German Space Agency DLR with funds provided by the Federal Ministry of Economics and Technology (BMWi) under grant number DLR 50WM1131.

SYQG 2.3 Tue 15:15 E 415

Sensing single remote nuclear spins in Nitrogen-Vacancy centers — ●JAN HONERT¹, NAN ZHAO¹, BERNHARD SCHMID¹, MICHAEL KLAS¹, JUNICHI ISOYA², MATTHEW MARKHAM³, DANIEL TWITCHEN³, FEDOR JELEZKO⁴, REN-BAO LIU⁵, HELMUT FEDDER¹, and JÖRG WRACHTRUP¹ — ¹3. Physikalisches Institut, University Stuttgart, Pfaffenwaldring 57, 70569 Stuttgart, Germany — ²Graduate School of Library, Information and Media Studies, University of Tsukuba, 1-2 Kasuga, Tsukuba, Ibaraki 305-8550, Japan — ³Element Six Ltd, Ascot SL5 8BP, Berks, England — ⁴Institut für Quantenoptik, Universität Ulm, 89081 Ulm, Germany — ⁵Department of Physics and Centre for Quantum Coherence, The Chinese University of Hong Kong, Shatin,

New Territories, Hong Kong, China

The detection of single nuclear spins would be useful for fields ranging from basic science to quantum information technology. In addition, the ability to address weakly coupled nuclear spins in the solid state expands the number of addressable qubits surrounding the detector spin significantly. Here, we present the detection and identification of single and remote ¹³C nuclear spins embedded in nuclear spin baths surrounding a single electron spin of a nitrogen-vacancy centre in diamond. We are able to amplify and detect the weak magnetic field noise (~ 10 nT) from a single nuclear spin located about 3 nm from the centre using dynamical decoupling control, and achieve a detectable hyperfine coupling strength as weak as ~ 300 Hz. We also confirm the quantum nature of the coupling present the first steps and results towards manipulating those spins.

SYQG 2.4 Tue 15:30 E 415

A satellite based quantum test of Einstein's equivalence principle — ●CHRISTIAN SCHUBERT¹ and THE STE-QUEST ATI TEAM² — ¹Institut für Quantenoptik, LU Hannover — ²European Consortium

STE-QUEST [1] aims for performing a quantum test of Einstein's Equivalence principle by verifying the universality of the free propagation of matter waves on a satellite. A dual species atom interferometer will measure the differential acceleration of Bose-Einstein condensates of ⁸⁷Rb and ⁸⁵Rb. This is assumed to be zero if the inertial mass coincides with the gravitational mass. The Eötvös ratio derived from the differential signal will be determined with an accuracy of parts in 10^{15} beyond state-of-the-art precision of 10^{-13} established by lunar laser ranging and torsion balances.

The matter waves will be simultaneously prepared and interrogated with a free evolution time of 10 s enabled by the weightlessness conditions in space. Within a single cycle of 20 s a shot noise limited sensitivity to accelerations of $3 \cdot 10^{-12}$ m/s² is anticipated. The simultaneous interferometry is carried out in a double diffraction Mach-Zehnder geometry and allows for high suppression ratios of noise and bias terms.

In the talk the measurement principle will be presented, an overview of the preliminary payload design will be given, and the estimated error budget will be discussed.

STE-QUEST is a proposal for an M3 mission in the frame of the Cosmic Vision program of ESA.

[1] <http://sci.esa.int/ste-quest>

SYQG 2.5 Tue 15:45 E 415

General relativistic effects in quantum interference of "clocks" — ●MAGDALENA ZYCH¹, FABIO COSTA¹, IGOR PIKOVSKI¹, CASLAV BRUKNER¹, and TIMOTHY C. RALPH² — ¹Universität Wien — ²University of Queensland

Quantum mechanics and general relativity have been extensively and independently confirmed in many experiments. However, all experiments that measured the influence of gravity on quantum systems are still fully consistent with non-relativistic, Newtonian gravity. Here we discuss a novel effect in quantum interference experiments that can probe the interplay between quantum mechanics and general relativity.

We consider interference of a "clock" – a particle with some evolving degrees of freedom – placed in a superposition of two different gravitational potentials. According to general relativity each amplitude of the superposition will experience a different gravitational time dilation. Due to quantum complementarity the visibility of quantum interference will thus drop to the extent to which the information about the location becomes available from the "clock". The clock can be implemented in an internal degree of freedom of a massive particle or in the position of a photon. The proposed experiment would thus provide the first test of quantum mechanics in curved background.