

HL 17: Focus Session: Topological Phenomena in Synthetic Matter (joint session DS/HL)

Topological insulators are a striking example of materials in which topological invariants are manifested in robustness against perturbations. Topology has emerged as an abstract, yet surprisingly powerful, new paradigm for controlling the flow of an excitation, e.g. the flow of electrons or light. This interdisciplinary Focus Session aims at discussing the latest experimental and theoretical results in the fast developing field of topological phenomena in synthetic matter. The recent merging of topology and cold atoms, photonics, mechanics and many more fields promises a considerable impact on these disciplines. We bring together leading theoretical and experimental experts from the fields of topological phenomena in synthetic matter to discuss recent progress and interdisciplinary synergy emerging at the interface of these fields. Furthermore, we give an overview to young scientists of exciting possibilities of interdisciplinary research in these fields with the special focus on the practical applications of fundamental science.

Organizer: Sebastian Klemmt (Julius-Maximilians-Universität Würzburg)

Time: Thursday 13:30–16:15

Location: H1

Topical Talk HL 17.1 Thu 13:30 H1
Exceptional Topology of Non-Hermitian Systems: from Theoretical Foundations to Novel Quantum Sensors — ●JAN CARL BUDICH — Institute of Theoretical Physics, TU Dresden, Dresden, Germany

In a broad variety of physical settings ranging from classical metamaterials to open quantum systems, non-Hermitian (NH) Hamiltonians have proven to be a powerful and conceptually simple tool for effectively describing dissipation. Motivated by recent experimental discoveries, investigating the topological properties of such NH systems has become a major focus of current research. In this talk, I give a brief introduction to this rapidly growing field, and present our latest results. Specifically, we discuss the occurrence of novel topological phases unique to NH systems. There, the role of spectral degeneracies familiar from Hermitian systems such as Weyl semimetals is played by exceptional points at which the effective NH Hamiltonian becomes non-diagonalizable. Furthermore, we show how guiding principles of topological matter such as the bulk boundary correspondence are qualitatively changed in the NH realm. Finally, we demonstrate that the sensitivity of NH systems to small changes in the boundary conditions may be harnessed to devise novel high-precision sensors.

Topical Talk HL 17.2 Thu 14:00 H1
In situ fabrication of (Bi,Sb)-based topological insulator - superconductor hybrid devices — ●PETER SCHÜFFELGEN — Forschungszentrum Jülich

With their experimental verification in 2007, topological insulators render a new and fascinating material class. A band inversion in the bulk of a 3D topological insulator creates a 2D metallic Dirac system at the physical surface of those 3D crystals. The surface Dirac states are topologically protected and have their spin locked to their momentum. This intrinsic quantum spin texture promises to enable fundamentally new, yet elusive quantum technologies, such as Majorana quantum bits. In this talk, I will introduce the material class of (Bi,Sb)-based topological insulators and discuss experimental challenges. I will present an in situ process that makes it possible to construct hybrid devices comprised of topological and superconductive nanostructures fully under ultra-high vacuum conditions via molecular beam epitaxy. A combi-

nation of stencil lithography and selective area growth allows for the realization of a variety of superconductor-topological insulator hybrid devices and solves the associated fabrication challenges.

Topical Talk HL 17.3 Thu 14:30 H1
Atomic monolayers as two-dimensional topological insulators — ●RALPH CLAESSEN — Physikalisches Institut und Würzburg-Dresden Cluster of Excellence ct.qmat, Universität Würzburg, Germany

Two-dimensional topological insulators (2D-TIs) are characterized by hosting spin-polarized conducting band states at their one-dimensional (1D) edges, giving rise to the quantum spin Hall (QSH) effect. As pointed out in the seminal work of Kane and Mele, graphene would constitute the most simple realization of a QSH insulator if it were not for its almost negligible spin-orbit interaction. It has been suggested that going to heavier group IV monolayers (such as the Sn-derived "stanene") could remedy this problem, but a convincing demonstration of such 2D TIs is still lacking. Recently we discovered that the neighboring groups III and V in the Periodic Table provide a promising alternative. Here I will discuss rational design, epitaxial synthesis, as well as ARPES and STM studies of two such synthetic QSH insulators, namely Bi (bismuthene) and In (indenene) monolayers grown on SiC(0001) substrates.

15 minutes break

Topical Talk HL 17.4 Thu 15:15 H1
Topological Insulator Lasers — ●MORDECHAI SEGEV — Technion - Israel Institute of Technology

Topological Insulator Lasers are semiconductor emitters fabricated on a potential landscape designed to harness the features of topological insulators to force injection-locking of the emitters, making them act as a single coherent laser. The concepts underlying topological insulator lasers will be reviewed along with the recent progress.

Topical Talk HL 17.5 Thu 15:45 H1
TBA — ●MORAIS SMITH — TBA
 TBA