

HK 49: Structure and Dynamics of Nuclei IX

Time: Wednesday 17:30–19:00

Location: SCH/A118

HK 49.1 Wed 17:30 SCH/A118

Measurements of the reaction cross sections of neutron-rich Sn isotopes at R³B setup. — ●ELEONORA KUDAIBERGENOVA¹, THOMAS AUMANN^{1,2,4}, MARTINA FEIJOO FONTAN⁵, ANDREA HORVAT^{1,3}, IVANA LIHTAR³, VALERII PANIN², and DOMINIC ROSSI^{1,2} for the R3B-Collaboration — ¹Institut für Kernphysik, TU Darmstadt, Germany — ²GSI Helmholtzzentrum für Schwerionenforschung, Darmstadt, Germany — ³Rudjer Boskovic Institute, Zagreb, Croatia — ⁴Helmholtz Forschungszentrum für Energie, Darmstadt, Germany — ⁵IGFAE, Universidad de Santiago de Compostela, Spain

The equation of state (EoS) is fundamental for understanding the structure of nuclear matter. The study of asymmetric nuclear matter via properties of neutron-rich nuclei became a current focus of investigation. The asymmetry term of the nuclear EoS is expressed by the symmetry energy at saturation J and its slope L , which has not yet been constrained well experimentally. It has been identified that a precise determination of the neutron removal cross section of neutron-rich nuclei, which is directly related to the neutron skin, would provide a much better constraint on L . To this end, an experiment was performed with the neutron-rich tin isotopes in the mass range $A=124-134$ on ¹²C targets at the R³B setup at the GSI/FAIR facility in inverse kinematics with very large acceptance. In this report, the current detector calibration, analysis status is presented.

This project was supported by the BMBF project No. 05P21RDFN2, the Helmholtz Research Academy Hessen for FAIR, and the GSI-TU Darmstadt cooperation.

HK 49.2 Wed 17:45 SCH/A118

Investigation of γ -softness: Lifetime measurements in ^{104,106}Ru — ●ARWIN ESMAYLZADEH¹, ANDREY BLAZHEV¹, KOSUKE NOMURA², JAN JOLIE¹, MARCEL BECKERS¹, CHRISTOPH FRANSEN¹, ROSA-BELLE GERST¹, ANDREAS HARTER¹, VASIL KARAYONCHEV^{1,3}, LUKAS KNAFLA¹, MARIO LEY¹, and FRANZISKUS VON SPEE¹ — ¹Institute for Nuclear Physics, University of Cologne — ²Department of Physics, University of Zagreb — ³TRIUMF, Canada

Lifetimes of the 2_1^+ , 4_1^+ , 6_1^+ , 2_2^+ and 3_2^+ states in ^{104,106}Ru were measured using the recoil distance Doppler shift technique and the Cologne Plunger device. Low-lying excited states in both nuclei were populated in a ¹⁰⁴Ru(¹⁸O, ¹⁸O)¹⁰⁴Ru* inelastic scattering and in a ¹⁰⁴Ru(¹⁸O, ¹⁶O)¹⁰⁶Ru two-neutron transfer reaction using the Cologne FN Tandem accelerator. The experimental energy levels and deduced electromagnetic transition probabilities are compared in the context of γ -softness and the mapped interacting boson model with input from the microscopic self-consistent mean-field calculation using a Gogny interaction [1]. The newly obtained results for the γ band, give a more detailed insight about the triaxial behavior of ^{104,106}Ru. The results will be discussed in the context of γ soft and rigid triaxial behavior which is present in the neutron-rich Ru isotopes [2]. This work supported by BMBF erbundprojekt 05P2021 (ErUM-FSP T07) grant 05P21PKFN1.

[1] K. Nomura et al., Phys. Rev. C 94, 044314 (2016)

[2] A. Esmaylzadeh et al., Phys. Rev. C (accepted in PRC) (2022)

HK 49.3 Wed 18:00 SCH/A118

Lifetime measurement of neutron rich Xe isotopes applying Fast-Timing method — ●ANDI MESSINGSCHLAGER¹, MARTIN VON TRESCKOW¹, THORSTEN KRÖLL¹, MATTHIAS RUDIGIER¹, ANDREY BLAZHEV², JULIA FISCHER², SORIN PASCU³, and JONATHAN N. WILSON⁴ for the nu-Ball2 N-SI-120-Collaboration — ¹TU Darmstadt — ²U Cologne — ³U Surrey — ⁴IJCLab Orsay

¹⁴²Xe is a neutron rich even-even isotope which lies between the double shell closure $N = 82$ and $Z = 50$ and a region in which an increased quadrupole and octupole collectivity is expected [1,2]. The lifetimes of excited states of ¹⁴²Xe are located in the range of some picoseconds. In this time range the Fast-Timing method is suited to determine the lifetime of excited states. Since there are different results for the lifetimes of excited states of the Xe-isotopes in experiments using the Fast-Timing method [1] and Coulomb excitation [2]. Therefore, we are going to analyse the data taken following the fission of ²³⁸U induced by a pulsed neutron beam of 1.7 MeV energy from the LICORNE neutron source. The nu-Ball2 multidetector array consisted of 24 HPGe Clover

detectors and 20 LaBr₃(Ce) detectors which promise excellent energy and time resolution, respectively. The campaign was performed at IJCLab in Orsay, France. Preliminary results will be presented. Supported by BMBF under Verbundprojekt 05P2021 (ErUM-FSP T07) grant 05P21RDFN1 and ARIEL

[1] S. Ilieva et al., PRC 94, 034302 (2016).

[2] C. Henrich, Dissertation TU Darmstadt (2020)

HK 49.4 Wed 18:15 SCH/A118

Gamma-ray spectroscopy of neutron-rich ^{55,57,59}Sc isotopes — ●RADOSTINA ZIDAROVA¹, MARTHA LILIANA CORTÉS², VOLKER WERNER¹, PAVLOS KOSEOGLOU¹, NORBERT PIETRALLA¹, PIETER DOORNENBAL², and ALEXANDRE OBERTELLI¹ — ¹TU Darmstadt, Germany — ²RIKEN-RIBF, Japan

Experimental data have shown that far from the valley of stability new magic numbers can emerge and the traditional ones can disappear. In particular, two new magic numbers at $N=32$ and $N=34$ have been suggested in the vicinity of $Z=20$ based on gamma-ray spectroscopy and mass measurements. In order to assess the impact of a single valence proton outside of the $Z=20$ shell on the shell-evolution mechanism in this region, it is necessary to study the neutron-rich Sc isotopes around, and even beyond, neutron number $N=34$. Investigation of exotic nuclei in this region was the goal of the third SEASTAR campaign at RIKEN-RIBF. Neutron-rich isotopes in the vicinity of ⁵³K were produced by fragmentation of a primary ⁷⁰Zn beam on a ⁹Be target. Known and new γ -ray transitions of the isotope ⁵⁵Sc were observed and new γ -rays from ^{57,59}Sc identified for the first time. Observed γ spectra from ^{55,57,59}Sc will be presented together with preliminary level schemes. They will be discussed in the framework of the tensor-driven shell evolution.

Supported by BMBF under Grant Nos. 05P19/21RDFN1.

HK 49.5 Wed 18:30 SCH/A118

Lifetime measurements of excited states in ⁵⁷Mn — ●HANNAH KLEIS, PETER REITER, KONRAD ARNSWALD, MAXIMILIAN DROSTE, ANDREY BLAZHEV, RAMONA BURGGRAF, and CRISTOPH FRANSEN — Institut für Kernphysik, Universität zu Köln

Previously, the $N = 32$ subshell closure was observed in the even-even Ca-, Ti-, and Cr-isotopes [1]. Adding more valence protons to the $\pi(f_{7/2})$ orbital reduces the shell gap at $N = 32$ which vanishes completely at ⁵⁸Fe. Lifetime measurements in the odd-even ⁵⁷Mn nucleus were performed in order to close the gap between $Z = 24$ and $Z = 26$. Excited states of ⁵⁷Mn were populated via ⁵⁵Mn(¹⁸O, ¹⁶O)⁵⁷Mn two-neutron transfer reactions at a beam energy of 38 MeV employing the FN tandem accelerator at the University of Cologne. The Doppler-shift attenuation method is utilized to determine new lifetimes for the $11/2_1^-$ and $9/2_1^-$ states. The experimentally determined transition probabilities are confronted with results from the GXPF1A shell-model interaction along the Mn-isotopes. The experimental findings in ⁵⁷Mn are well reproduced by this interaction. The comparison of excitation energies and $B(E2)$ strengths is extended to all odd-even nuclei between Ca and Ni with neutron numbers $N = 26$ and $N = 36$ in order to discuss the nature of the $N = 32$ subshell closure.

[1] D. Steppenbeck et al., Nature 502, 7470 (2013)

HK 49.6 Wed 18:45 SCH/A118

Lifetime measurements of neutron-rich Kr isotopes within the nu-Ball2 fission campaign — ●J. FISCHER¹, A. BLAZHEV¹, C. HIVER², J. JOLIE¹, A. MESSINGSCHLAGER³, S. PASCU⁴, M. VON TRESCKOW³, N. WARR¹, and J. N. WILSON² for the nu-Ball2 N-SI-120-Collaboration — ¹U Cologne — ²IJCLab Orsay — ³TU Darmstadt — ⁴U Surrey

Nuclei beyond the band of stability are crucial to our understanding of the atomic nucleus and nuclear forces. In recent years, neutron-rich krypton isotopes have been studied as part of various campaigns. New gamma-transitions and levels were discovered in the first nu-Ball and SEASTAR-2015 campaigns [1,2] which compared to theory indicated oblate-prolate shape coexistence already in ⁹⁶Kr [2]. However, the limited information on transition strengths did not allow for firm conclusions. Therefore lifetime measurements were performed at the IJCLab Orsay as part of the nu-Ball2 fission campaign. The nuclei of interest were produced with a fast-neutron-induced fission reac-

tion $^{238}\text{U}(n,f)$. The improved multidetector-array (nu-Ball2), a novel hybrid γ -spectrometer consisting of HPGe and LaBr3(Ce) detectors provided excellent energy and timing information, respectively. The fast-timing method allows for lifetime determination down to about 10 ps and thus a possibility to determine transition strengths in the nu-

clei of interest. Preliminary results will be presented. *Supported by BMBF under Verbundprojekt 05P2021 (ErUM-FSP T07) grant 05P21PKFN1. / [1] R.-B. Gerst et al., PRC 102, 064323 (2020). ; [2] R.-B. Gerst et al., PRC 105, 024302 (2022).